

Package leaflet: Information for the patient

Olopatadine 1 mg/mL eye drops, solution

Olopatadine

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Olopatadine is and what it is used for
2. What you need to know before you use Olopatadine
3. How to use Olopatadine
4. Possible side effects
5. How to store Olopatadine
6. Contents of the pack and other information

1. What Olopatadine is and what it is used for

Olopatadine is a medicine for treatment of allergic conditions of the eye. It works by reducing the intensity of the allergic reaction.

Olopatadine is used for the treatment of signs and symptoms of seasonal allergic conjunctivitis.

Allergic conjunctivitis. Some materials (allergens) like pollens, house dust or animal fur may cause allergic reactions resulting in itching, redness as well as swelling of the surface of your eye.

2. What you need to know before you use Olopatadine

Do not use Olopatadine

- If you are allergic (hypersensitive) to olopatadine or any of the other ingredients of this medicine (listed in section 6).
- You should not use Olopatadine if you are breast-feeding.

Warnings and precautions

Talk to your doctor or pharmacist before using Olopatadine.

You should remove contact lenses that are in your eyes before using Olopatadine.

Children

Do not use Olopatadine in children under the age of 3 years. Do not give this medicine to children under the ages of 3 years because there is no data to indicate that it is safe and work in children under 3 years.

Other medicines and Olopatadine

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

If you are using other eye drops or eye ointment medicines, leave at least 5 minutes between each medicine. Eye ointments should be administered last.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advise before using this medicine.

You should not use Olopatadine if you are breast-feeding, ask your doctor for advise before using this medicine.

Driving and using machines

You may find that your vision is blurred for a time just after you use Olopatadine. Do not drive or use machines until this has worn off.

Olopatadine contains Benzalkonium chloride

Benzalkonium chloride may cause eye irritation and is known to discolour soft contact lenses, therefore contact with soft contact lenses should be avoided. If you wear contact lens you should remove contact lenses prior to application and wait at least 15 minutes before putting your lenses back in.

3. How to use Olopatadine

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Dosage

The recommended dose is one drop in the eye or eyes, twice a day – morning and evening.

Use this much unless your doctor tells you to do differently. Only use Olopatadine in both eyes if your doctor told you to. Use it for as long as your doctor told you to.

Olopatadine should only be used as an eye drop.

Method of administration




Fig. 1

Fig. 2

- Get the Olopatadine bottle and a mirror.
- Wash your hands.
- Take the bottle and twist off the cap.
- Hold the bottle, pointing down, between your thumb and middle finger.
- Tilt your head back. Pull down your eyelid with a clean finger, until there is a 'pocket' between the eyelid and your eye. The drop will go in here (picture 1).
- Bring the bottle tip close to the eye. Use the mirror if it helps.

- Don't touch your eye or eyelid, surrounding areas or other surfaces with the dropper. It could infect the drops left in the bottle.
- Gently press on the base of the bottle to release one drop of Olopatadine at a time.
- Don't squeeze the bottle, it is designed so that just a gentle press on the bottom is needed (picture 2).
- If you use drops in both eyes, repeat the steps for your other eye.
- Put the bottle cap back on firmly immediately after use.

If a drop misses your eye, try again.

If you use more Olopatadine than you should

Rinse it all out with warm water. Do not put in any more drops until it's time for your next regular dose.

If you forget to use Olopatadine

Use a single drop as soon as you remember, and then go back to your regular dosing routine.

However, if it is almost time for your next dose, skip the missed dose before going back to your regular dosing routine. Do not use a double dose to make up for the one missed.

If you stop using Olopatadine

Do not stop using this medicine without first speaking to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects have been seen with olopatadine:

Common: may affect up to 1 in 10 people

Effects in the eye: eye pain, eye irritation, dry eye, abnormal eye sensation, eye discomfort.

General side effects: headache, fatigue, dry nose, bad taste.

Uncommon: may affect up to 1 in 100 people

Effects in the eye: blurred, reduced, or abnormal vision, corneal disorder, eye surface inflammation with or without surface damage, inflammation or infection of the conjunctiva, eye discharge, sensitivity to light, increased tear production, itchy eye, redness of the eye, eyelid abnormality, itching, redness, swelling, or crusting of the eyelid.

General side effects: abnormal or decreased sensation, dizziness, runny nose, dry skin, skin inflammation.

Not known: frequency cannot be estimated from the available data

Effects in the eye: eye swelling, corneal swelling, change in pupil size.

General side effects: shortness of breath, increased allergic symptoms, facial swelling, drowsiness, generalized weakness, nausea, vomiting, sinus infection, skin redness and itching.

In very rare cases, some patients with severe damage to the clear layer at the front of the eye (the cornea) have developed cloudy patches on the cornea due to calcium build-up during treatment.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in

Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Olopatadine

Keep out of reach of children.
At a temperature below 25°C.
Do not freeze.
Store in a tightly closed bottle.

Do not use this medicine after the expiry date which is stated on the bottle and the box after 'Exp'.
The expiry date refers to the last day of the month.
Shelf-life after first opening: 28 days at a temperature below 25°C.

Do not throw away medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Content of the pack and other information

What Olopatadine contains

- The active substance is olopatadine. Each mL of solution contains 1 mg of olopatadine (as hydrochloride).
- The other ingredients are benzalkonium chloride, disodium phosphate dihydrate (E339), sodium chloride, hydrochloric acid (E507) and/or sodium hydroxide, water for injections.

What Olopatadine looks like and contents of the pack

Olopatadine is a clear and colourless solution, free from visible particles.
The finished product is packed in white LDPE bottles of 5 ml with natural nozzle and white screw cap with tamper-evident ring.
One bottle in a carton box.

Marketing Authorisation Holder

Antibiotic-Razgrad AD
68, Aprilsko vastanie Blvd., office 201
Razgrad 7200
Bulgaria

Manufacturer

Balkanpharma-Razgrad AD
68 "Aprilsko vastanie" Blvd.
7200 Razgrad
Bulgaria

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder.

Antibiotic-Razgrad AD

68, Aprilsko vastanie Blvd., office 201
Razgrad 7200
Bulgaria
Tel: +359 (84) 660 999
Fax: +359 (84) 634 272
e-mail: office@antibiotic.bg

This leaflet was last revised in 06/2018